

Domain decomposition methods for nonlinear transmission conditions

Matthias A. F. Gsell*, Olaf Steinbach

Institut für Numerische Mathematik, TU Graz,
Steyrergasse 30, 8010 Graz, Austria

Abstract

We consider the transformation of quasilinear partial differential equations to a coupled system of linear equations, but with nonlinear transmission conditions on the interfaces. After deriving a variational formulation, we will discuss Mortar finite element discretization strategies and present a numerical example.

1 Model problem

As a model problem we consider the following quasilinear boundary value problem in a bounded domain $\Omega \subset \mathbb{R}^d$ ($d = 2, 3$) with Lipschitz boundary $\Gamma = \partial\Omega = \bar{\Gamma}_D \cup \bar{\Gamma}_N$ and $\Gamma_D \cap \Gamma_N = \emptyset$ to find p such that

$$\left. \begin{aligned} -\nabla \cdot (k_r(\theta(p)) \nabla p) &= f && \text{in } \Omega, \\ p &= 0 && \text{on } \Gamma_D, \\ k_r(\theta(p)) \nabla p \cdot \mathbf{n} &= g_N && \text{on } \Gamma_N. \end{aligned} \right\} \quad (1)$$

This model problem can be seen as a simplification of the stationary Richards equation without gravity, see [1].

If k_r only depends on θ , we can introduce

$$u(\mathbf{x}) := \kappa(p(\mathbf{x})) := \int_0^{p(\mathbf{x})} k_r(\theta(s)) \, ds$$

as the Kirchhoff transformation of p . Here we assume $k_r \circ \theta$ to be Lipschitz and bounded with

$$0 < c_l \leq k_r(\theta(\cdot)) \leq c_u < \infty.$$

Therefore we obtain

$$\nabla u = \kappa'(p) \nabla p = k_r(\theta(p)) \nabla p$$

and problem (1) can be transformed to the following linear problem to find u such that

$$-\Delta u = f \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \Gamma_D, \quad \nabla u \cdot \mathbf{n} = g_N \quad \text{on } \Gamma_N.$$




Figure 1: several domains




Figure 2: dual interface mesh

Obviously, such a simplification can not be achieved if k_r depends in addition explicitly on $\mathbf{x} \in \Omega$. However, if this dependency is piecewise within non-overlapping subdomains Ω_i , we can introduce local Kirchhoff transformations $u_i(\mathbf{x}) := \kappa_i(p_i(\mathbf{x}))$ for $\mathbf{x} \in \Omega_i$. So we can reformulate problem (1) and obtain the following nonlinear transmission problem, in the case of two subdomains, see Figure 1, to find u_i , $i = 1, 2$, such that

$$\left. \begin{aligned} -\Delta u_i &= f && \text{in } \Omega_i, \\ u_i &= 0 && \text{on } \Gamma_{D,i}, \\ \nabla u_i \cdot \mathbf{n}_i &= g_{N,i} && \text{on } \Gamma_{N,i}, \\ \nabla u_1 \cdot \mathbf{n} = \nabla u_2 \cdot \mathbf{n}, \quad \kappa_1^{-1}(u_1) &= \kappa_2^{-1}(u_2) && \text{on } \Gamma_I, \end{aligned} \right\} \quad (2)$$

where $\Gamma_I = \partial\Omega_1 \cap \partial\Omega_2$, $\Gamma_{D,i} = \partial\Omega_i \cap \Gamma_D$, $\Gamma_{N,i} = \partial\Omega_i \cap \Gamma_N$ and $g_{N,i} = g_N|_{\Gamma_{N,i}}$. The second, nonlinear interface condition ensures the continuity of the solution p of the original problem (1), where we started from. Next we will derive a variational formulation for the nonlinear transmission problem (2).

2 Primal–hybrid formulation

In [2,3], an equivalent primal–hybrid formulation for the Poisson equation is presented. The idea is to introduce Lagrange multipliers acting on interfaces which ensure global continuity of the solution. We use this approach for the nonlinear transmission problem (2) and obtain the following variational problem to find $(u, \lambda) \in X \times M$, such that

$$\left. \begin{aligned} \sum_{i=1}^2 \int_{\Omega_i} \nabla u_i \cdot \nabla v_i \, dx + \int_{\Gamma_I} \lambda [v]_{\Gamma_I} \, ds_x \\ = \sum_{i=1}^2 \int_{\Omega_i} f_i v_i \, dx + \sum_{i=1}^2 \int_{\Gamma_{N,i}} g_{N,i} v_i \, ds_x \quad \text{for all } v \in X, \\ \int_{\Gamma_I} \mu (u)_{\Gamma_I} \, ds_x = 0 \quad \text{for all } \mu \in M \end{aligned} \right\} \quad (3)$$

with the ansatz spaces

$$X := \{v \in L_2(\Omega) \mid v_i = v|_{\Omega_i} \in H_{0,\Gamma_{D,i}}^1(\Omega_i), i = 1, 2\}, \quad M := H_{00}^{\frac{1}{2}}(\Gamma_I)',$$

and with the jump terms

$$[v]_{\Gamma_I} := (v_1 - v_2)|_{\Gamma_I} \in M' \quad \text{and} \quad (u)_{\Gamma_I} := (\kappa_1^{-1}(u_1) - \kappa_2^{-1}(u_2))|_{\Gamma_I} \in M'.$$

*Corresponding author, e-mail gsell@tugraz.at

3 Discretization

For the discretization of (3) we introduce the finite dimensional subspaces

$$X_h := \{v \in L_2(\Omega) \mid v_i \in S_h^1(\mathcal{T}_{h,i}), i = 1, 2\} \subset X$$

and

$$M_h := \{\mu \in L_2(\Gamma_I) \mid \mu \in S_h^0(\mathcal{I}'_h)\} \subset M,$$

where $\mathcal{T}_{h,i}$ is an admissible triangulation of Ω_i , and \mathcal{I}'_h is a modified dual mesh along the interface Γ_I as depicted in Figure 2, see also [4]. For the solution of the nonlinear discrete problem we apply Newton's method by solving a sequence of the following linear problems. For given data $w_h \in X_h$, find $(u_h, \lambda_h) \in X_h \times M_h$, such that

$$\left. \begin{aligned} & \sum_{i=1}^2 \int_{\Omega_i} \nabla u_{h,i} \cdot \nabla v_{h,i} \, dx + \int_{\Gamma_I} \lambda_h [v_h]_{\Gamma_I} \, ds_x \\ & = \sum_{i=1}^2 \int_{\Omega_i} f_i v_{h,i} \, dx + \sum_{i=1}^2 \int_{\Gamma_{N,i}} g_{N,i} v_{h,i} \, ds_x \quad \forall v_h \in X_h, \\ & \int_{\Gamma_I} \mu_h (w_h, u_h)'_{\Gamma_I} \, ds_x \\ & = \int_{\Gamma_I} \mu_h (w_h, w_h)'_{\Gamma_I} \, ds_x - \int_{\Gamma_I} \mu_h (w_h)_{\Gamma_I} \, ds_x \quad \forall \mu \in M_h \end{aligned} \right\} \quad (4)$$

where

$$(w_h, u_h)'_{\Gamma_I} := ((\kappa_1^{-1})'(w_{h,1}) u_{h,1} - (\kappa_2^{-1})'(w_{h,2}) u_{h,2})_{\Gamma_I}$$

is the linearization of the nonlinear jump $(u_h)_{\Gamma_I}$ and w_h corresponds to the previous Newton iteration.

4 Numerical example

Consider $\mathbb{R}^2 \supset \bar{\Omega} = [0, 1]^2 = ([0, 0.5] \times [0, 1]) \cup ([0.5, 1] \times [0, 1]) =: \bar{\Omega}_1 \cup \bar{\Omega}_2$, where $\Gamma_N = \{0\} \times (0, 1)$ and $\Gamma_D = \partial\Omega \setminus \Gamma_N$. The local diffusion coefficients are chosen as in [1, Chapter 1]. The remaining data $f, g_{N,i}, g_{D,i}$ are given appropriately.




Figure 3: (discontinuous) solution u of problem (4) and its (continuous) inverse Kirchhoff transformations $p_i = \kappa_i^{-1}(u_i)$, $i = 1, 2$.

Acknowledgement: This work was supported by the Austrian Science Fund (FWF) within the IGDK 1754.

References

- [1] H. BERNINGER, Domain Decomposition Methods for Elliptic Problems with Jumping Nonlinearities and Application to the Richards Equation, PhD thesis, Freie Universität Berlin, 2008.
- [2] D. BOFFI, F. BREZZI, and M. FORTIN, Mixed finite element methods and applications, Springer, Heidelberg, 2013.
- [3] P. A. RAVIART and J. M. THOMAS, Primal hybrid finite element methods for 2nd order elliptic equations, *Math. Comp.* **31**(138), 391–413 (1977).
- [4] B. I. WOHLMUTH, Discretization methods and iterative solvers based on domain decomposition, Springer, 2001.